CORNELL UNIVERSITY

Department of Physics

Physics 214 Prelim I Fall 2003

NAME:

SECTION:

Instructions

— To receive credit, you <u>must</u> place your answers in the boxes provided whenever required.

— Closed book; no notes. You may use a calculator.

— Check that you have all $\underline{14}$ pages (including cover page). The formula sheet is distributed separately.

— Important note: Except for some challenge problems, each part of this exam is designed to be answered without the answers of previous parts. The parts within a given problem become more and more difficult. If you get stuck on one part, skip to the next problem and come back later if you have more time.

Problem	Score	Grader
1. (16 pts)		
2. (20 pts)		
3. (24 pts)		
4. (12 pts)		
5. (12 pts)		
6. (16 pts)		
Total (100 pts)		

Contents

1	Problem 1: Standing Waves on Strings(a) Sketches (12 points)(b) Comparing frequencies (4 points)	$\begin{bmatrix} 16 \text{ points} \end{bmatrix}$	3 3 3
2	Problem 2: Lab Experiment I (a) Speed of sound in aluminum (8 points) (b) Bulk modulus of aluminum (8 points) (c) Spring constant of the rod (4 points) <i>Challenge Problem!</i>	[20 points]	4 4 5
3	Problem 3: Harmonic Oscillator with Friction(a) Equation of Motion (6 points)	[24 points]	6 6 7 8 8 9
4	Problem 4: Equation of <i>transverse</i> motion in a fluid	[12 points]	10
5	Problem 5: Sound Waves versus Waves on a String(a) Equation for strings (8 points)	[12 points]	12 12 12
6	Problem 6: Generalized Boundary Conditions for a Sound Tube(a) Air Pressure and Displacement of the Piston (6 points)(b) Equation of Motion for the piston (6 points)(c) Standing wave solutions (4 points) <i>Challenge Problem!</i>	[16 points]	13 13 14 14

List of Figures

1	Wave patterns for different boundary conditions.	3
2	A mass-spring realization of a harmonic oscillator with sliding friction.	6
3	A chunk of fluid experiencing viscous forces	10
4	Generalized boundary conditions for a sound tube.	13



Figure 1: Wave patterns for different boundary conditions.

Problem 1: Standing Waves on Strings [16 points] 1

(a)Sketches (12 points)

On the axes which Figure 1 provides, sketch the form of the lowest frequency mode for each of the three possible combinations of boundary conditions (fixed-fixed, fixed-free, free-free) for a string. For full credit, label all points where y = 0 and all points where $\partial y / \partial x = 0$.

Comparing frequencies (4 points) (b)

Among the three possible combinations of boundary conditions from part (a), which allows the lowest possible standing-wave frequency for the same length L, tension τ and mass M of the string?

Provide your answer in the box below.



2 Problem 2: Lab Experiment I [20 points]

As part of the first laboratory experiment in Phys 214, you measured the frequency of the lowest mode of an aluminum rod of length L = 1 m. You should have found a frequency near $f = 2 \times 10^3$ Hz. (For the purpose of this problem, take this to be the exact value.) Because you held the rod at its center while leaving both ends free, this gives the frequency corresponding to the free-free combination of boundary conditions from Problem 3. Finally, further measurements on the actual rod you used show it to have a cross sectional area $A = \pi r^2 = 500 \text{ mm}^2 = 500 \times 10^{-6} \text{ m}^2$ and a total mass M = 1.35 kg.

(a) Speed of sound in aluminum (8 points)

Use the above information to compute the speed of sound in the rod. Give a numerical result in units of m/s.

(b) Bulk modulus of aluminum (8 points)

The actual speed of sound in Aluminum is c = 5100 m/s. (Note: You will not find this value from the data in part (a) because of the fake value given there for f!) Using the actual speed of sound c = 5100 m/s and the information given above, compute the bulk modulus B of aluminum. Give a numerical result in units of N/m^2 .

(c) Spring constant of the rod (4 points) Challenge Problem!

The rod can be used as a very stiff spring. When the length of the rod is changed by an amount Δx , the rod pushes back with a restoring force $F = -k \Delta x$. Use the definition

$$\Delta P = -B\frac{\Delta V}{V_0},$$

to compute a numerical value for the proportionality constant k in units of N/m. Hint: For this problem, you may ignore any tendency for the radius of the rod to increase as the rod is compressed. The influence of this effect is small in aluminum.



Figure 2: A mass-spring realization of a harmonic oscillator with sliding friction.

3 Problem 3: Harmonic Oscillator with Friction[24 points]

Consider a harmonic oscillator with mass m acted on by an ideal spring of spring constant k and equilibrium position $x = x_{eq} = 0$ and by a sliding friction force in the direction opposite to the motion and of magnitude $f = \mu mg$ where μ is the coefficient of sliding friction and g the acceleration of gravity. (See Figure 2.)

(a) Equation of Motion (6 points)

Show that, when the mass moves to the right, the Equation of Motion for the harmonic oscillator with friction is:

$$\frac{d^2x}{dt^2} + \mu g + \omega_0^2 x = 0 , \qquad \omega_0 \equiv \sqrt{\frac{k}{m}} .$$
 (3.1)

(b) General solution (6 points)

Show that the following is a general solution to the equation of motion (3.1):

$$x(t) = -\frac{\mu g}{\omega_0^2} + \Re \mathfrak{e} \left[\underline{A} e^{i\omega_0 t}\right].$$
(3.2)

Note: Be sure to list *each* requirement for a general solution as you check it.

(c) Real Part of \underline{A} (4 points)

For parts (c)-(e), consider initial conditions where, at t = 0, the mass is at the equilibrium position $x_0 = x_{eq} = 0$ and moving with a velocity $v_0 > 0$ to the right.

Find the real part $\Re [\underline{A}]$ given the above initial conditions and general solution. Express your answer in terms of only v_0 , μ , g, and ω_0 . (Note: you may not need all of these.)

(d) Imaginary Part of \underline{A} (4 points)

Find the imaginary part $\Im [\underline{A}]$ under the same conditions. Express your answer in terms of only v_0 , μ , g, and ω_0 . (Note: you may not need all of these.)

(e) Maximum distance (4 points) Challenge Problem!

What maximum position x_{max} will the mass reach before stopping and turning around? Express your answer in terms of only v_0 , μ , g, and ω_0 . (Note: you may not need all of these.)



Figure 3: A chunk of fluid experiencing viscous forces

4 Problem 4: Equation of *transverse* motion in a fluid [12 points]

Consider transverse motion along the y direction of a fluid with density ρ_0 and bulk modulus B. For plane waves propagating along x, the constitutive relation for such motion is that the driving force per unit area on a chunk at x due to the adjacent fluid is

$$\frac{F_y^{\text{visc}}(x,t)}{A} = c_v \frac{\partial^2 y(x,t)}{\partial x \,\partial t},\tag{4.1}$$

with the direction of the force as in Figure 3. Here, c_v is a constant known as the coefficient of viscosity (a constant which is large for "thick" fluids like honey).

Which of the following formulas best represents the equation for transverse motion in a fluid? (Provide your answer in the box on the next page.)

$$\begin{aligned} \text{(A)} \quad c_v \frac{\partial^3 y(x,t)}{\partial x^2 \, \partial t} &= \rho_0 \frac{\partial^2 y(x,t)}{\partial t^2} \\ \text{(B)} \quad c_v \frac{\partial^2 y(x,t)}{\partial x \, \partial t} &= \rho_0 \frac{\partial^2 y(x,t)}{\partial t^2} \\ \text{(C)} \quad B \frac{\partial^2 y(x,t)}{\partial x^2} &= \rho_0 \frac{\partial^2 y(x,t)}{\partial t^2} \\ \text{(D)} \quad c_v \frac{\partial^3 y(x,t)}{\partial x \, \partial t^2} + B \frac{\partial^2 y(x,t)}{\partial x^2} &= \rho_0 \frac{\partial^2 y(x,t)}{\partial t^2} \\ \text{(E)} \quad c_v \frac{\partial^3 y(x,t)}{\partial x^2 \, \partial t} + B \frac{\partial^2 y(x,t)}{\partial x^2} &= \rho_0 \frac{\partial^2 y(x,t)}{\partial t^2} \end{aligned}$$

PLEASE TURN PAGE

PLEASE TURN PAGE

5 Problem 5: Sound Waves versus Waves on a String [12 points]

The following equation is true for sound waves:

$$-\frac{\partial}{\partial x}\left[xB\frac{\partial s}{\partial x} - Bs\right] + \frac{\partial}{\partial t}\left[x\rho_0\frac{\partial s}{\partial t}\right] = 0.$$
(5.1)

(a) Equation for strings (8 points)

What is the analogous equation for strings?

(b) Another conservation law (4 points) Challenge Problem!

Conservation of what physical quantity does your equation for strings represent? Hint: The density of the conserved quantity appears inside the $\frac{\partial}{\partial t}[$], and the driving "force" for the conserved quantity appears inside the $\frac{\partial}{\partial x}[$]. PSfrag replacements



Figure 4: Generalized boundary conditions for a sound tube.

6 Problem 6: Generalized Boundary Conditions for a Sound Tube [16 points]

A tube of length L is filled with air of bulk modulus B and mass density ρ_0 at atmospheric pressure P_0 . One end of the tube (x = L) is closed. At the other end (x = 0), a massless piston of area A can slide freely (without friction) along the tube. The piston is attached to a *shock absorber* which generates a horizontal external force on the piston $\vec{F}^{(\text{ext})} = -b\vec{v}$, where \vec{v} is the velocity of the piston. (See Figure 4.) We denote the sound displacement inside the tube as s(x,t) and the pressure inside the tube as P(x,t).

(a) Air Pressure and Displacement of the Piston (6 points)

Assuming that the piston is moving to the right with velocity $v = \partial s(x = 0)/\partial t$, draw a free body diagram for the piston, indicating the directions and the magnitudes of all the forces using only P(x = 0, t), $\partial s(x = 0, t)/\partial t$, b, A, ρ_0 , L, and P_0 . (Note: you may not need all of these.)

(b) Equation of Motion for the piston (6 points)

Assuming small amplitude waves, so that you can take the location of the piston to be $x \approx 0$, write the Equation of Motion for the piston in terms of only the constants describing the problem $(b, A, B, \rho_0, L, \text{ and } P_0)$, and s(x, t) and its derivatives evaluated at x = 0. (Note: you may not need all of these.)

(c) Standing wave solutions (4 points) Challenge Problem!

In the limit $b \to \infty$, what is the frequency of the lowest frequency standing sound wave in the tube *in terms of* L, ρ_0 , and B?